

From ODLRO to the Meissner Effect and Flux Quantization

Chi Au and Bao-Heng Zhao*

Department of Applied Mathematics

Hong Kong Polytechnic University, Hong Kong

February 1, 2008

Abstract

It has been shown that the electron system with ODLRO in the reduced density matrix ρ_2 can not support a uniform magnetic field, i.e., ODLRO in ρ_2 implies the Meissner effect [2][3], furthermore, the magnetic field trapped in the system is quantized[3]. This note extends above results in two cases. We show that (1) the system with ODLRO in ρ_2 can not support a non uniform, cylindrically symmetric magnetic field; (2) the system with ODLRO in ρ_2 can not support a magnetic field slowly varying in space, and the magnetic flux trapped in it is quantized.

*Permanent address: Department of Physics, Graduate School, Chinese Academy of Sciences, P. O. Box 3908, Beijing 100039, China

PACS numbers: 05.30.-d; 74.20.-z; 4.25.Ha

Introduction

In 1962 C. N. Yang proposed that the superconductivity is characterized by the existence of off-diagonal long-range order (ODLRO) in the two-particle reduced density matrix ρ_2 for a fermion system[1]. However, there was no clear-cut proof to show directly that ODLRO in ρ_2 leads to superconductivity for a long time. In recent works [2][3], it has been proved that ODLRO in ρ_2 implies that a non vanishing uniform magnetic field $\mathbf{B}$ can not exist in an electron system, i.e., there is the Meissner effect. Furthermore, it has been shown that the magnetic flux trapped in the system is quantized[3]. However, the question of whether a non uniform magnetic field can be expelled from the system with ODLRO in ρ_2 is still open. The present work extends [2] and [3] in two cases. We show that 1. the system with ODLRO in ρ_2 can not support a non uniform, cylindrically symmetric magnetic field; 2. the system with ODLRO in ρ_2 can not support a non uniform magnetic field, which is slowly varying in space, and the magnetic flux trapped in the system is quantized.

System with Cylindric Symmetry

Consider the case that N electrons are in a circular cylinder. We chose a cylindric coordinates with z-axis along the central line of the cylinder. In which $\mathbf{r} = (R, \theta, z)$, and $\hat{\mathbf{R}}, \hat{\theta}$ and $\hat{\mathbf{z}}$ are unit vectors along the R, θ and z direction, respectively. Suppose that the magnetic field is cylindrically

symmetric, and the gauge to be [4]

$$\mathbf{A}(\mathbf{r}) = -\theta R B(R) \hat{\mathbf{R}}, \quad (1)$$

where $B(R)$ is a continuous function. From (1), we have $\mathbf{B}(\mathbf{r}) = \nabla \times \mathbf{A}(\mathbf{r}) = B(R) \hat{\mathbf{z}}$. Suppose that the system undergoes an infinitesimal rotation $\hat{\mathbf{z}} \delta\theta$,

$$\begin{aligned} \mathbf{A}(\mathbf{r}) &\rightarrow -(\theta - \delta\theta) R B(R) \hat{\mathbf{R}} = \mathbf{A}(\mathbf{r}) + \nabla \chi_{\delta\theta}(\mathbf{r}), \\ \chi_{\delta\theta}(\mathbf{r}) &= \delta\theta \int_{R_0}^R R' B(R') dR', \end{aligned} \quad (2)$$

where R_0 is a constant. Eq.(2) shows that the rotation induces a gauge transformation. We shall see that it enable us to show that $\mathbf{B}(\mathbf{r}) = 0$ as the system has ODLRO in the two particle reduced density matrix ρ_2 .

The Schrödinger equation of the electrons is

$$\begin{aligned} \frac{1}{2m} \sum_j \left[\frac{\hbar}{i} \nabla_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j) \right]^2 \psi_n(\mathbf{r}_1, \dots, \mathbf{r}_N) + U(\mathbf{r}_1, \dots, \mathbf{r}_N) \psi_n(\mathbf{r}_1, \dots, \mathbf{r}_N) \\ = E_n \psi_n(\mathbf{r}_1, \dots, \mathbf{r}_N). \end{aligned} \quad (3)$$

We assume that the potential $U(\mathbf{r}_1, \dots, \mathbf{r}_N)$ depends only on the distances between electrons. Under the rotation, (3) becomes

$$\begin{aligned} \frac{1}{2m} \sum_j \left[\frac{\hbar}{i} \nabla_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j) \right]^2 \psi'_n(\mathbf{r}_1, \dots, \mathbf{r}_N) \\ + U(\mathbf{r}_1, \dots, \mathbf{r}_N) \psi'_n(\mathbf{r}_1, \dots, \mathbf{r}_N) = E_n \psi'_n(\mathbf{r}_1, \dots, \mathbf{r}_N), \end{aligned} \quad (4)$$

where

$$\begin{aligned} \psi'_n(\mathbf{r}_1, \dots, \mathbf{r}_N) &= \psi_n(\mathbf{r}_1 - \delta\mathbf{r}_1, \dots, \mathbf{r}_N - \delta\mathbf{r}_N) \exp\left[\frac{ie}{\hbar c} \sum_j \chi_{\delta\theta}(\mathbf{r}_j)\right], \\ \delta\mathbf{r}_i &= \hat{\theta}_i R_i \delta\theta. \end{aligned} \quad (5)$$

Both $\{\psi_n\}$ and $\{\psi'_n\}$ are complete sets of orthonormal eigenfunctions of the Hamiltonian, which are single-valued in $\mathbf{r}_i$.

The element of the reduced density matrix for two particles can be expressed as

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \int \cdots \int \frac{d\mathbf{r}_3 \cdots d\mathbf{r}_N}{(N-2)!} \frac{1}{Z} \sum_n \exp(-E_n/kT) \\ &\times \psi_n(\mathbf{r}'_1, \mathbf{r}'_2, \mathbf{r}_3, \cdots, \mathbf{r}_N) \psi_n^*(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \cdots, \mathbf{r}_N), \end{aligned} \quad (6)$$

where Z is the partition function, and T the temperature. Substituting ψ'_n for ψ_n makes no change for ρ_2 , since ρ_2 is basis independent. So we also have

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \int \cdots \int \frac{d\mathbf{r}_3 \cdots d\mathbf{r}_N}{(N-2)!} \frac{1}{Z} \sum_n \exp(-E_n/kT) \\ &\times \psi'_n(\mathbf{r}'_1, \mathbf{r}'_2, \mathbf{r}_3, \cdots, \mathbf{r}_N) \psi'^*_n(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \cdots, \mathbf{r}_N), \end{aligned} \quad (7)$$

Expressing the primed ψ'_n in (7) in terms of the non primed ψ_n , we obtain

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \exp\left\{\frac{ie}{c\hbar}[\chi_{\delta\theta}(\mathbf{r}'_1) + \chi_{\delta\theta}(\mathbf{r}'_2) - \chi_{\delta\theta}(\mathbf{r}_1) - \chi_{\delta\theta}(\mathbf{r}_2)]\right\} \\ &\times \int \cdots \int \frac{d\mathbf{r}_3 \cdots d\mathbf{r}_N}{(N-2)!} \frac{1}{Z} \sum_n \exp(-E_n/kT) \\ &\times \psi_n(\mathbf{r}'_1 - \delta\mathbf{r}'_1, \mathbf{r}'_2 - \delta\mathbf{r}'_2, \mathbf{r}_3 - \delta\mathbf{r}_3, \cdots, \mathbf{r}_N - \delta\mathbf{r}_N) \\ &\times \psi_n^*(\mathbf{r}_1 - \delta\mathbf{r}_1, \mathbf{r}_2 - \delta\mathbf{r}_2, \mathbf{r}_3 - \delta\mathbf{r}_3, \cdots, \mathbf{r}_N - \delta\mathbf{r}_N). \end{aligned} \quad (8)$$

Shift the integration variables, $\theta_i \rightarrow \theta_i + \delta\theta$, $i = 3, \dots, N$. Note that $\mathbf{r}_i - \delta\mathbf{r}_i = (R_i, \theta_i - \delta\theta, z_i)$. We obtain a relation for ρ_2 at different space points:

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \exp\left\{\frac{ie}{c\hbar}[\chi_{\delta\theta}(\mathbf{r}'_1) + \chi_{\delta\theta}(\mathbf{r}'_2) - \chi_{\delta\theta}(\mathbf{r}_1) - \chi_{\delta\theta}(\mathbf{r}_2)]\right\} \\ &\times \rho_2(\mathbf{r}'_1 - \delta\mathbf{r}'_1, \mathbf{r}'_2 - \delta\mathbf{r}'_2; \mathbf{r}_1 - \delta\mathbf{r}_1, \mathbf{r}_2 - \delta\mathbf{r}_2). \end{aligned} \quad (9)$$

By definition, the ODLRO in ρ_2 means that

$$\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) \rightarrow \Phi(\mathbf{r}'_1, \mathbf{r}'_2) \Phi^*(\mathbf{r}_1, \mathbf{r}_2), \quad (10)$$

as $|\mathbf{r}'_i - \mathbf{r}_j|$, $i, j = 1, 2$, approaches macroscopic distances, and $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ does not vanish as $|\mathbf{r}_1 - \mathbf{r}_2|$ and $|\mathbf{r}'_1 - \mathbf{r}'_2|$ remain finite in microscopic scale. Where Φ is the eigenfunction of ρ_2 with the largest eigenvalue. If we assume ODLRO in ρ_2 , (8) and (9) lead to

$$\begin{aligned} \Phi(\mathbf{r}_1, \mathbf{r}_2) &= \exp[i\alpha(\delta\theta)] \exp\left\{\frac{ie}{c\hbar}[\chi_{\delta\theta}(\mathbf{r}_1) + \chi_{\delta\theta}(\mathbf{r}_2)]\right\} \\ &\times \Phi(\mathbf{r}_1 - \delta\mathbf{r}_1, \mathbf{r}_2 - \delta\mathbf{r}_2), \end{aligned} \quad (11)$$

where $\alpha(\delta\theta)$ is a real function of $\delta\theta$, and it is independent of $\mathbf{r}_1, \mathbf{r}_2$. It is easily seen that $\alpha(\delta\theta + \delta\theta') = \alpha(\delta\theta) + \alpha(\delta\theta')$, $\alpha(0) = 0$.

Rotating the system successively, as $\sum \delta\theta = 2\pi$, we have $\sum \delta\mathbf{r}_1 = \sum \delta\mathbf{r}_2 = 0$. $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ is single-valued because ψ_n is single-valued. Thus, we obtain from (10)

$$\exp[i\alpha(2\pi)] \exp\left\{\frac{ie}{c\hbar}2\pi\left[\int_{R_0}^{R_1} RB(R)dR + \int_{R_0}^{R_2} RB(R)dR\right]\right\} = 1. \quad (12)$$

(12) leads to

$$\frac{c\hbar}{e}\alpha(2\pi) + 2\pi\left[\int_{R_0}^{R_1} RB(R)dR + \int_{R_0}^{R_2} RB(R)dR\right] = n\frac{ch}{e}, \quad n = \text{integers}. \quad (13)$$

On the left hand side of (13), R_1 and R_2 can be changed continuously. Let $R_1 \rightarrow R_1 + dR_1$, the left hand of (13) gets an infinitesimal increment $2\pi R_1 B(R_1) dR_1$. On the other hand, the right hand side only takes discrete values. To be consistent, we have to take $B(R_1) = 0$. Since R_1 is arbitrary in the system, we obtain

$$\mathbf{B}(\mathbf{r}) = 0. \quad (14)$$

Thus a non vanishing cylindrically symmetric magnetic field can not exist in the system with ODLRO in ρ_2 . In other words, the magnetic field is expelled from the system, i.e., there is the Meissner effect.

Electrons in the Magnetic Field Slowly Varying in Space

Consider the case of electrons in a magnetic field which is slowly varying in space. The vector potential can be written as

$$\mathbf{A}(\mathbf{r}) = \mathbf{A}_0(\mathbf{r}) + \nabla\varphi(\mathbf{r}), \quad (15)$$

and $\mathbf{B}(\mathbf{r}) = \nabla \times \mathbf{A}_0(\mathbf{r})$. In general, $\varphi(\mathbf{r})$ is multi-valued, however, $\nabla\varphi(\mathbf{r})$ is single-valued.

Under an infinitesimal translation $\delta\mathbf{r}$,

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}_0(\mathbf{r} - \delta\mathbf{r}) + \nabla\varphi(\mathbf{r} - \delta\mathbf{r}) = \mathbf{A}(\mathbf{r}) - \delta\mathbf{r} \cdot \nabla[\mathbf{A}_0(\mathbf{r}) + \nabla\varphi(\mathbf{r})]. \quad (16)$$

In (16)

$$\begin{aligned} \delta\mathbf{r} \cdot \nabla \mathbf{A}_0 &= \nabla(\delta\mathbf{r} \cdot \mathbf{A}_0) - \delta\mathbf{r} \times \mathbf{B} \\ &= \nabla(\delta\mathbf{r} \cdot \mathbf{A}_0) - \nabla[\delta\mathbf{r} \cdot (\mathbf{B} \times \mathbf{r})] + [(\delta\mathbf{r} \cdot \nabla)\mathbf{B}] \times \mathbf{r} + \delta\mathbf{r} \times [(\mathbf{r} \cdot \nabla)\mathbf{B}]. \end{aligned} \quad (17)$$

Provided that $\mathbf{B}(\mathbf{r})$ is slowly varying, such that

$$|\delta\mathbf{r} \times \mathbf{B}| \gg |[(\delta\mathbf{r} \cdot \nabla)\mathbf{B}] \times \mathbf{r} + \delta\mathbf{r} \times [(\mathbf{r} \cdot \nabla)\mathbf{B}]| \quad (18)$$

i.e., roughly speaking, $|\nabla B_i| / |\mathbf{B}|$, $i = x, y, z$, is much less than L^{-1} , where L is the linear dimension of the system. Under this condition, we can write

$$\delta\mathbf{r} \cdot \nabla \mathbf{A}_0 = \nabla(\delta\mathbf{r} \cdot \mathbf{A}_0) - \nabla[\delta\mathbf{r} \cdot (\mathbf{B} \times \mathbf{r})], \quad (19)$$

and (16) becomes

$$\mathbf{A}(\mathbf{r}) \rightarrow \mathbf{A}(\mathbf{r}) + \nabla[\delta\mathbf{r} \cdot (\mathbf{B} \times \mathbf{r}) - \delta\mathbf{r} \cdot \mathbf{A}_0 - \delta\mathbf{r} \cdot \nabla\varphi(\mathbf{r})]. \quad (20)$$

We see that the infinitesimal translation induces a gauge transformation.

In the Hamiltonian of the system,

$$H = \sum_j \frac{1}{2m} [-i\hbar \nabla_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j)]^2 + U(\mathbf{r}_1, \dots, \mathbf{r}_N), \quad (21)$$

we assume that the potential $U(\mathbf{r}_1, \dots, \mathbf{r}_N)$ depends only on the distances between electrons. Suppose that $\{\psi_n\}$ is a complete set of single-valued orthonormal eigenfunctions of the Hamiltonian. Then $\{\psi'_n\}$ is also a complete set of single-valued orthonormal eigenfunctions of the Hamiltonian[3], where

$$\psi'_n(\mathbf{r}_1, \dots, \mathbf{r}_N) = e^{\frac{ie}{c\hbar} \sum_j \chi_{\delta\mathbf{r}}(\mathbf{r}_j)} \psi_n(\mathbf{r}_1 - \delta\mathbf{r}, \dots, \mathbf{r}_N - \delta\mathbf{r}), \quad (22)$$

and

$$\chi_{\delta\mathbf{r}}(\mathbf{r}_j) = \delta\mathbf{r} \cdot [\mathbf{B}(\mathbf{r}_j) \times \mathbf{r}_j - \mathbf{A}_0(\mathbf{r}_j) - \nabla_j \varphi(\mathbf{r}_j)]. \quad (23)$$

Repeating the discussion similar to that leads to (8) (see also [3]), we have

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \exp\left\{\frac{ie}{c\hbar} [\chi_{\delta\mathbf{r}}(\mathbf{r}'_1) + \chi_{\delta\mathbf{r}}(\mathbf{r}'_2) - \chi_{\delta\mathbf{r}}(\mathbf{r}_1) - \chi_{\delta\mathbf{r}}(\mathbf{r}_2)]\right\} \\ &\times \rho_2(\mathbf{r}'_1 - \delta\mathbf{r}, \mathbf{r}'_2 - \delta\mathbf{r}; \mathbf{r}_1 - \delta\mathbf{r}, \mathbf{r}_2 - \delta\mathbf{r}). \end{aligned} \quad (24)$$

If there is ODLRO in ρ_2 , (10) is valid, and

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) = \exp[i\alpha(\delta\mathbf{r})] \exp\left\{\frac{ie}{c\hbar} [\chi_{\delta\mathbf{r}}(\mathbf{r}_1) + \chi_{\delta\mathbf{r}}(\mathbf{r}_2)]\right\} \Phi(\mathbf{r}_1 - \delta\mathbf{r}, \mathbf{r}_2 - \delta\mathbf{r}), \quad (25)$$

where $\alpha(\delta\mathbf{r})$ is real, and not dependent on $\mathbf{r}_1, \mathbf{r}_2$. It is easily seen that $\alpha(\delta\mathbf{r}_1 + \delta\mathbf{r}_2) = \alpha(\delta\mathbf{r}_1) + \alpha(\delta\mathbf{r}_2)$, and $\alpha(0) = 0$. Repeating successively for infinitesimal displacements along a closed path, and noting the single-valuedness of $\Phi(\mathbf{r}_1, \mathbf{r}_2)$, we obtain from (25) the relation

$$\exp\left\{i \sum \alpha(\delta\mathbf{r})\right\} \exp\left\{\frac{2ie}{c\hbar} \oint d\mathbf{r} \cdot [\mathbf{B}(\mathbf{r}) \times \mathbf{r} - \mathbf{A}_0(\mathbf{r}) - \nabla\varphi(\mathbf{r})]\right\} = 1, \quad (26)$$

where the summation $\sum \alpha(\delta \mathbf{r})$ is over the closed path. In a simply connected region, around a sufficiently small contour, on which $\mathbf{B}$ can be considered as a constant vector, we have $\oint d\mathbf{r} \cdot [\mathbf{B}(\mathbf{r}) \times \mathbf{r}] = 2\phi$, $\oint d\mathbf{r} \cdot \mathbf{A}_0(\mathbf{r}) = \phi$, where ϕ is the magnetic flux through the surface spanned by the small contour, and $\oint d\mathbf{r} \cdot \nabla \varphi(\mathbf{r}) = 0$, $\sum \alpha(\delta \mathbf{r}) = 0$. We obtain,

$$\exp\left\{\frac{2ie}{c\hbar} \oint d\mathbf{r} \cdot [\mathbf{B}(\mathbf{r}) \times \mathbf{r} - \mathbf{A}_0(\mathbf{r})]\right\} = \exp\left\{\frac{2ie}{c\hbar} \phi\right\} = 1. \quad (27)$$

Thus

$$\phi = n \frac{ch}{2e}, \quad n = \text{integers}. \quad (28)$$

The contour can be deformed continuously. If $\mathbf{B} \neq 0$, the flux ϕ can be changed continuously. Which is in contradiction to the right hand side of (28). To be consistent, we conclude that $\mathbf{B} = 0$. It means that the system with ODLRO in ρ_2 can not support a magnetic field slowly varying in space, i.e., there is the Meissner effect.

Since $\mathbf{B} = 0$, we can take $\mathbf{A}_0 = 0$. When the path is not simply connected, e.g., it winds around an inaccessible tunnel, the path can not be deformed across this inaccessible region. In this case we still have $\sum \alpha(\delta \mathbf{r}) = 0$ [5], from (26), we obtain $\exp\left\{\frac{2ie}{c\hbar} \oint d\mathbf{r} \cdot \nabla \varphi(\mathbf{r})\right\} = 1$, thus

$$\oint d\mathbf{r} \cdot \nabla \varphi(\mathbf{r}) = n \frac{ch}{2e}, \quad n = \text{integers}. \quad (29)$$

This equation shows that the magnetic flux trapped in the tunnel is quantized.

In summary, we have shown that if there is ODLRO in the reduced density matrix ρ_2 , 1. the system can not support a cylindrically symmetric magnetic field; 2. the system can not support a slowly varying magnetic field. In these

two cases, we have derived the Meissner effect from the existence of ODLRO in ρ_2 . Furthermore, in the latter case, we have shown that the magnetic flux trapped in the system is quantized. We know that the magnetic field is not vanishing, e.g., in the boundary region in superconductor. This fact is not in contradiction to our result, since the non vanishing magnetic field is in the region where it varies rapidly.

Acknowledgment

This work is supported in part by The Hong Kong Polytechnic University. One of the authors (B. H. Zhao) is grateful to the hospitality of the Department of Applied Mathematics, The Hong Kong Polytechnic University. He is also supported in part by NNSF of China.

References

- [1] C. N. Yang, Rev. Mod. Phys. **34**, 694(1962).
- [2] G. L. Swell, J. Stat. Phys. **61**, 415(1990).
- [3] H. T. Nieh, G. Su and B. H. Zhao, Off-diagonal long range order: Meissner effect and flux quantization (Preprint, 1994), to appear in Phys. Rev. **B**.
- [4] M. Bawin and A. Burnel, J. Phys. A **16**, 2173(1983); F. Wilczek, Phys. Rev. Lett. **48**, 1144(1982); **49**, 957(1982).
- [5] For a detail discussion about $\exp\{i\sum\alpha(\delta\mathbf{r})\} = 1$ on a closed path, see [3], where $\exp\{i\sum\alpha(\delta\mathbf{r})\}$ is written as $F(C)$.